CONSENT FORM

for

UROLOGICAL SURGERY

(Designed in compliance with



Department consent form 1)

Name of proposed procedure (include brief explanation if medical term not clear)	ANAESTHETIC
RADICAL CYSTECTOMY AND ILEAL CONDUIT (MALE)	μμ ⁻ GENERAL/REGIONAL
THIS INVOLVES REMOVAL OF ENTIRE BLADDER, PROSTATE AND PELVIC LYMPH NODES AND	μμ ⁻ LOCAL
PERMANENT DIVERSION OF URINE TO ABDOMINAL SKIN USING A LOOP OF BOWEL AS A STOMA	μμ ⁻ S EDATION

Serious or frequently occurring risks

COMMON

- TEMORARY INSERTION OF A NASAL TUBE, DRAIN, STENT
- HIGH RISK OF IMPOTENCE (LACK OF ERECTIONS)
- DRY ORGASM WITH NO SEMEN PRODUCED CAUSING INFERTILITY
- BLOOD TRANSFUSION REQUIRED

OCCASIONAL

- NEED TO REMOVE THE PENILE URINARY PIPE AS PART OF THE PROCEDURE
- BLOOD LOSS REQUIRING REPEAT SURGERY
- CANCER MAY NOT BE CURED WITH REMOVAL OF BLADDER ALONE

RARE

- INFECTION OR HERNIA OF INCISION REQUIRING FURTHER TREATMENT
- ANAESTHETIC OR CARDIOVASCULAR PROBLEMS POSSIBLY REQUIRING INTENSIVE CARE ADMISSION (INCLUDING
- CHEST INFECTION, PULMONARY EMBOLUS, STROKE, DEEP VEIN THROMBOSIS, HEART ATTACK AND DEATH.)
- DECREASE RENAL FUNCTION WITH TIME

VERY RARE

- DIARRHOEA DUE TO SHORTENED BOWEL / VITAMIN DEFICIENCY REQUIRING TREATMENT
- BOWEL AND URINE LEAKAGE FROM ANASTOMOSIS REQUIRING RE-OPERATION
- SCARRING TO BOWEL OR URETERS REQUIRING OPERATION IN FUTURE
- SCARRING, NARROWING OR HERNIA FORMATION AROUND STOMAL OPENING REQUIRING REVISION
- INTRAOPERATIVE RECTAL INJURY REQUIRING COLOSTOMY

ALTERNATIVE TREATMENT: RADIATION TREATMENT TO BLADDER, CONTINENT DIVERSION OF URINE WAS DISCUSSED.