CONSENT FORM

for

UROLOGICAL SURGERY

(Designed in compliance with



Department consent form 1)

Name of proposed procedure (include brief explanation if medical term not clear)	ANAESTHETIC
RADICAL CYSTECTOMY AND FORMATION OF NEW BLADDER WITH BOWEL (FEMALE)	μμ [−] GENERAL/REGIONAL
THIS INVOLVES REMOVAL OF BLADDER, PELVIC LYMPH NODES & REMAINING FEMALE ORGANS (I.E.	μμμ [−] LOCAL
OVARIES AND UTERUS INCLUDING A PORTION OF VAGINA) AND FORMATION OF A NEW BLADDER	μμμ [−] S EDATION
USING BOWEL	μμμ [−] S

Serious or frequently occurring risks

COMMON

- TEMORARY INSERTION OF A NASAL TUBE, DRAIN, STENT
- PAIN OR DIFFICULTY WITH SEXUAL INTERCOURSE DUE TO NARROWING OR SHORTENING OF VAGINA
- NEED TO SELF CATHETERISE IF NEW BLADDER FAILS TO FULLY EMPTY
- IN EVENT OF OVARY REMOVAL MENOPAUSE MAY OCCUR
- BLOOD TRANSFUSION REQUIRED

OCCASIONAL

- BLOOD LOSS REQUIRING REPEAT SURGERY
- CANCER MAY NOT BE CURED WITH REMOVAL OF BLADDER ALONE
- INCONTINENCE OF URINE

RARE

- INFECTION OR HERNIA OF INCISION REQUIRING FURTHER TREATMENT
- ANAESTHETIC OR CARDIOVASCULAR PROBLEMS POSSIBLY REQUIRING INTENSIVE CARE ADMISSION (INCLUDING
- CHEST INFECTION, PULMONARY EMBOLUS, STROKE, DEEP VEIN THROMBOSIS, HEART ATTACK AND DEATH.)
- DECREASE RENAL FUNCTION WITH TIME

VERY RARE

- DIARRHOEA DUE TO SHORTENED BOWEL / VITAMIN DEFICIENCY REQUIRING TREATMENT
- BOWEL AND URINE LEAKAGE FROM ANASTOMOSIS REQUIRING RE-OPERATION
- SCARRING TO BOWEL OR URETERS REQUIRING OPERATION IN FUTURE
- URETHRAL RECURRENCE OF THE CANCER
- INTRAOPERATIVE RECTAL INJURY REQUIRING COLOSTOMY

ALTERNATIVE TREATMENT: RADIATION TREATMENT TO BLADDER.